

Theta line bundles and the determinant of the Hodge bundle.

Alexis Kouvidakis
 University of Crete
 Department of Mathematics
 71409, Iraklion-Crete, Greece
 e-mail: kouvid@math.uch.gr

Abstract: In this note we examine the question of expressing the determinant of the push forward of a symmetric line bundle on an abelian fibration in terms of the pull back of the relative dualizing sheaf via the zero section.

0 Introduction.

Let $f : X \longrightarrow B$ be a fibration of abelian varieties with a zero section $s : B \longrightarrow X$. Let $\mathcal{L}$ be a symmetric line bundle on X , trivialized along the zero section, which defines a polarization of type $D = (d_1, \dots, d_g)$ on the fibration. A theorem of Faltings-Chai, see Theorem 5.1 in Ch. 1 of [3], gives the following expression of the determinant of the push forward of the line bundle $\mathcal{L}$ in terms of the pull back $s^*\omega_{X/B}$ of the relative dualizing sheaf of the family:

$$8d^3 \det f_* \mathcal{L} = -4d^4 s^* \omega_{X/B},$$

where $d := d_1 \cdots d_g$.

In this note we show that in the complex analytic category, the above torsion factor can be improved. More specifically, we have

Theorem A *Let $f : X \longrightarrow B$ be a fibration of complex abelian varieties of relative dimension g and s its zero section. Let $\mathcal{L}$ be a symmetric line bundle on X , trivialized along the zero section, which defines a polarization of type $D = (d_1, \dots, d_g)$, where $d_1 | \dots | d_g$ are positive integers. Let $d = d_1 \cdots d_g$, then*

$$8 \det f_* \mathcal{L} = -4ds^* \omega_{X/B},$$

except when $3|d_g$ and $\gcd(3, d_{g-1}) = 1$, in which case we have that

$$24 \det f_* \mathcal{L} = -12ds^* \omega_{X/B}.$$

Moreover, when $\mathcal{L}$ is totally symmetric i.e. $\mathcal{L}$ is fiberwise the square of a symmetric line bundle, we have

Theorem B *Let $f : X \longrightarrow B$ be a fibration of complex abelian varieties of relative dimension $g \geq 3$ and s its zero section. Let $\mathcal{L}$ be a totally symmetric line bundle on X , trivialized along the zero section, which defines a polarization of type $D = (d_1, \dots, d_g)$, where $d_1 | \dots | d_g$ are positive even integers. Let $d = d_1 \cdots d_g$, then*

$$\det f_* \mathcal{L} = -\frac{d}{2} s^* \omega_{X/B},$$

except when $3|d_g$ and $\gcd(3, d_{g-1}) = 1$, in which case we have that

$$3 \det f_* \mathcal{L} = -3\frac{d}{2} s^* \omega_{X/B}.$$

The theorems are proved by using a “higher rank” version of the Theta Transformation Formula, see Propositions 1.1 and 1.2, in order to construct transition functions for $\det f_*(\mathcal{L})$, see Lemma 2.1.

At the end, we apply Theorem B to the case of the universal Jacobian variety $f_{g-1} : \mathcal{J}^{g-1} \longrightarrow \mathcal{M}_g$. This is an abelian torsor which parametrizes line bundles of degree $g-1$ on the fibers of the universal curve, where $\mathcal{M}_g$ denotes the moduli space of smooth, irreducible curves of genus $g \geq 3$, without automorphisms. On that we have a canonical theta divisor defined by the push forward of the universal symmetric product of degree $g-1$, via the Abel-Jacobi map. We denote by Θ the corresponding line bundle and by λ the determinant of the Hodge bundle of the universal curve $\psi : \mathcal{C} \longrightarrow \mathcal{M}_g$, i.e. $\lambda = \det \psi_* \omega_{\mathcal{C}/\mathcal{M}_g}$. We then have

Theorem C *Keeping the above notation, we have that*

$$\det f_{g-1 *} (\Theta^{\otimes n}) = \frac{1}{2} n^g (n-1) \lambda.$$

We also give an alternative way of proving Theorem C by utilizing special properties of the universal Jacobian varieties.

Acknowledgment. I would like to thank Professor L. Moret-Bailly for showing me the alternative way for proving Theorem C.

1 Abelian varieties and theta functions.

We begin by recalling some standard theory for abelian varieties and theta functions. The main reference we follow is the book by Lange and Birkenhake [4]. We also keep (most of) the notation of that book. We denote by $X = V/\Lambda$ an abelian variety, V is a $\mathbb{C}$ -vector space of dimension g and Λ a $2g$ -full lattice in V .

1.1 Line bundles on abelian varieties.

On the abelian variety $X = V/\Lambda$, a line bundle is given, up to isomorphism, by data (H, χ) , where H is a hermitian form on V s.t. $\text{Im}H(\Lambda, \Lambda) \subset \mathbb{Z}$ and $\chi : \Lambda \rightarrow \mathbb{C}_1^*$ a semicharacter. The isomorphism class of line bundles on X given by the above data is denoted by $L(H, \chi)$, see Ch. 2, & 2 in [4].

Let $\phi : X' = V'/\Lambda' \rightarrow X = V/\Lambda$ be a map of abelian varieties. To that one can associate two maps, namely the analytic representation $\phi_a : V' \rightarrow V$ and the rational representation $\phi_r : \Lambda' \rightarrow \Lambda$, see Ch. 1, & 2 in [4]. If ϕ is a map of abelian varieties as above, then $\phi^*L(H, \chi) = L(\phi_a^*H, \phi_r^*\chi)$.

1.2 Decomposition of V - Characteristics.

Given a hermitian form H as in Section 1.1, there is a base $\langle \lambda_1, \dots, \lambda_g; \lambda_{g+1}, \dots, \lambda_{2g} \rangle$ of Λ , w.r.t. which $\text{Im}H$ is given by a matrix of the form $\begin{pmatrix} 0 & D \\ -D & 0 \end{pmatrix}$, where D is an integral diagonal matrix with elements $d_1 | \dots | d_g$. Such a base is called a symplectic base for H . When the matrix D can be chosen to be the identity, then H is called a principal polarization of X .

Let Λ_1 be the lattice spanned by the first g vectors and Λ_2 the lattice spanned by the last g vectors. If $\lambda \in \Lambda$, we write $\lambda = \lambda_1 + \lambda_2$ for the corresponding decomposition of λ . Then V , viewed as an $\mathbb{R}$ -vector space, decomposes as $V = V_1 \oplus V_2$, where $V_i = \Lambda_i \otimes \mathbb{R}$; we call that one decomposition of V for H . If $v \in V$, we write $v = v_1 + v_2$ for the corresponding decomposition of v . We denote by $E := \text{Im}H$ the alternating form associated to H and by B the symmetric form on V which is the $\mathbb{C}$ -bilinear extension of the symmetric form $H|_{V_2 \times V_2}$, see Lemma 2.1 of Ch. 3 in [4]. We define $\Lambda(H) := \{v \in V; \text{Im}H(v, \Lambda) \subset \mathbb{Z}\}$. Then $\Lambda(H) = \Lambda(H)_1 \oplus \Lambda(H)_2$, where $\Lambda(H)_i := \Lambda(H) \cap V_i$.

We denote by $\text{Pic}^H(X)$ the group of line bundles, up to isomorphism, corresponding to the Hermitian form H . After choosing a decomposition of V for H , we define in $\text{Pic}^H(X)$ a distinguished isomorphism class of line bundles, denoted by $L_0 \cong L(H, \chi_0)$, by setting $\chi_0(\lambda) =$

$e(\pi i E(\lambda_1, \lambda_2))$. If $L(H, \chi)$ is any other element of $\text{Pic}^H(X)$, then there exists a $c \in V$, unique up to translations by elements of $\Lambda(H)$, s.t. $L(H, \chi) = T_c^* L(H, \chi_0)$, where T_c is the translation map by c on X . Note that if $L(H, \chi)$ is of characteristic c then $\chi(\lambda) = \chi_0(\lambda) e(2\pi i E(c, \lambda))$, see Ch. 3, & 1 in [4].

1.3 Canonical - Classical factor of automorphy.

A factor of automorphy is a holomorphic map $f : \Lambda \times V \longrightarrow \mathbb{C}^\times$ satisfying the condition $f(\lambda_1 + \lambda_2, v) = f(\lambda_1, \lambda_2 + v) f(\lambda_2, v)$. Two factors of automorphy f and f' are called equivalent iff there is a holomorphic function $h : V \rightarrow \mathbb{C}^\times$ satisfying $f'(\lambda, v) = f(\lambda, v) h(v) h(v + \lambda)^{-1}$. We use the notation $f' = f \star h$ to indicate the above situation.

Let $L(H, \chi)$ be an isomorphism class of line bundles on X . We define the canonical factor of automorphy $a_{(H, \chi)} : \Lambda \times V \longrightarrow \mathbb{C}^\times$ by

$$a_{(H, \chi)}(\lambda, v) := \chi(\lambda) e(\pi H(v, \lambda) + \frac{\pi}{2} H(\lambda, \lambda)).$$

We define the classical factor of automorphy $e_{(H, \chi)} : \Lambda \times V \longrightarrow \mathbb{C}^\times$ by

$$e_{(H, \chi)}(\lambda, v) := \chi(\lambda) e(\pi(H - B)(v, \lambda) + \frac{\pi}{2}(H - B)(\lambda, \lambda)).$$

We have that $e_{(H, \chi)}(\lambda, v) = a_{(H, \chi)}(\lambda, v) e(\frac{\pi}{2} B(v, v)) e(\frac{\pi}{2} B(v + \lambda, v + \lambda))^{-1}$. This shows that the above factors are equivalent. When the semicharacter is the χ_0 i.e. the characteristic is 0, we simplify the notation for the above factors of automorphy to a_H and e_H .

1.4 Period matrices.

Let $X = V/\Lambda$ be an abelian variety and H a polarization of type D . We choose a symplectic base $\langle \lambda_1, \dots, \lambda_g; \lambda_{g+1}, \dots, \lambda_{2g} \rangle$ of Λ for H . Since $\langle \lambda_{g+1}, \dots, \lambda_{2g} \rangle$ is a $\mathbb{C}$ -base of V , there exists a unique $g \times 2g$ matrix Π of the form $\Pi = (Z, D)$, where $Z \in \mathfrak{h}_g$, with $\langle \lambda_1, \dots, \lambda_g \rangle = \langle \lambda_{g+1}, \dots, \lambda_{2g} \rangle Z$. We call Π the period matrix of X corresponding to the above choice of a symplectic base. Therefore to the triple $(X, H, \langle \lambda_1, \dots, \lambda_g; \lambda_{g+1}, \dots, \lambda_{2g} \rangle)$ corresponds an element $Z \in \mathfrak{h}_g$. Conversely, given $Z \in \mathfrak{h}_g$, let $\Lambda_Z := (Z, D)\mathbb{Z}^{2g}$. Define $X_Z := \mathbb{C}^g/\Lambda_Z$. Let H_Z be the Hermitian form with matrix $(\text{Im} Z)^{-1}$ w.r.t. the standard base of $\mathbb{C}^g$. This one defines then a polarization on X_Z of type D . Let λ_i be the vector corresponding to the i -column of the matrix (Z, D) . Then $\langle \lambda_1, \dots, \lambda_g; \lambda_{g+1}, \dots, \lambda_{2g} \rangle$ turns out to be a symplectic base for H_Z . Therefore to $Z \in \mathfrak{h}_g$ corresponds a triple $(X, H_Z, \langle \lambda_1, \dots, \lambda_g; \lambda_{g+1}, \dots, \lambda_{2g} \rangle)$ and the two constructions are inverse to each other.

1.5 Canonical - Classical theta functions.

Let $f : \Lambda \times V \longrightarrow \mathbb{C}^\times$ be a factor of automorphy. A theta function corresponding to f is a holomorphic function $\theta : V \longrightarrow \mathbb{C}$ satisfying the functional equation $\theta(\lambda + v) = f(\lambda, v)\theta(v)$. Let $L(H, \chi)$ be of characteristic c . Theta functions corresponding to the canonical (resp. classical) factor of automorphy $a_{(H, \chi)}$ (resp. $e_{(H, \chi)}$) are called canonical (resp. classical) theta functions for $L(H, \chi)$. After choosing a decomposition of V for H , we define

$$\theta_0^c(v) := e(-\pi H(v, c) - \frac{\pi}{2}H(c, c) + \frac{\pi}{2}B(v + c, v + c)) \sum_{\lambda_1 \in \Lambda_1} e(\pi(H - B)(v + c, \lambda_1) - \frac{\pi}{2}(H - B)(\lambda_1, \lambda_1)).$$

We have the following facts, see Ch. 3, & 2 in [4].

1. θ_0^c is a canonical theta function for $L(H, \chi)$. By the definition we deduce that $\theta_0^c(v) = e(-\pi H(v, c) - \frac{\pi}{2}H(c, c))\theta(v + c)$.
2. Let $\theta_w^c(v) := a_{(H, \chi)}(w, v)^{-1} \theta_0^c(v + w)$, where $\bar{w} \in K(H) := \Lambda(H)/\Lambda$. This is well defined and it is also a canonical theta function for $L(H, \chi)$.
3. The set $\langle \theta_w^c ; \bar{w} \in K(H)_1 := \Lambda(H)_1/\Lambda_1 \rangle$ forms a base of the canonical theta functions for $L(H, \chi)$.

Let $Z \in \mathfrak{h}_g$. We denote by j_Z the isomorphism of $\mathbb{R}$ -vector spaces $j_Z : \mathbb{R}^{2g} \longrightarrow \mathbb{C}^g$ which sends $x \mapsto (Z, 1)x$. Let Λ_D denote the lattice $\mathbb{Z}^g \oplus D\mathbb{Z}^g$ in $\mathbb{R}^{2g}$ and $\Lambda_Z := j_Z(\Lambda_D)$ the corresponding lattice in $\mathbb{C}^g$. Given $v \in \mathbb{C}^g$, we can write uniquely $v = Zv^1 + v^2$, with $v^i \in \mathbb{R}^g$. We then have that $\lambda \in \Lambda_Z$ can be written uniquely in the form $\lambda = Z\lambda^1 + \lambda^2$ with $\lambda^1 \in \mathbb{Z}^g$ and $\lambda^2 \in D\mathbb{Z}^g$.

Let now $L(H, \chi)$, where $H = H_Z$ is a polarization of type D , be an isomorphy class of line bundles of characteristic c . We have the following facts (most of them can be found in Ch. 8, & 5 in [4]; the rest is straight forward calculation):

1. $H(v, w) = {}^t v (\text{Im} Z)^{-1} \bar{w}$, $B(v, w) = {}^t v (\text{Im} Z)^{-1} w$.
 $(H - B)(v, w) = -2i {}^t v w^1$, $E(v, w) = {}^t v^1 w^2 - {}^t v^2 w^1$.
2. $e_H(\lambda, v) = e(-\pi i {}^t \lambda^1 Z \lambda^1 - 2\pi i {}^t v \lambda^1)$.
 $a_H(\lambda, v) = e(\pi i {}^t \lambda^1 \lambda^2 + \pi {}^t v (\text{Im} Z)^{-1} \bar{\lambda} + \frac{\pi}{2} {}^t \lambda (\text{Im} Z)^{-1} \bar{\lambda})$.
It is $e_H = a_H \star h$, with $h(v) = e(\frac{\pi}{2} {}^t v (\text{Im} Z)^{-1} v)$.

3. Let $\mathbb{Z}_D$ denote the group $\mathbb{Z}_D := \mathbb{Z}_{d_1} \oplus \dots \mathbb{Z}_{d_g}$.
 $\Lambda(H)_1 = \{Zv^1 \mid v^1 \in D^{-1}\mathbb{Z}^g\}$, $\Lambda(H)_2 = \{v^2 \mid v^2 \in \mathbb{Z}^g\}$.
 $K(H)_1 \cong D^{-1}\mathbb{Z}_D$, $K(H)_2 \cong \mathbb{Z}_D$.
4. Let $c = Zc^1 + c^2$. Then $\theta \begin{bmatrix} c^1 \\ c^2 \end{bmatrix} (v, Z) := e(-\frac{\pi}{2}B(v, v) + \pi i {}^t c^1 c^2) \theta_0^c(v)$ is a classical theta function for $L(H, \chi)$. We have
$$\theta \begin{bmatrix} c^1 \\ c^2 \end{bmatrix} (v, Z) = \sum_{\lambda^1 \in \mathbb{Z}^g} e\left(\pi i {}^t(\lambda^1 + c^1)Z(\lambda^1 + c^1) + 2\pi i {}^t(v + c^2)(\lambda^1 + c^1)\right).$$
5. Let $c = Zc^1 + c^2 \in \mathbb{C}^g$ and $w = Zw^1 + w^2 \in \Lambda(H)$. Then
 - a) $\theta \begin{bmatrix} c^1 + w^1 \\ c^2 \end{bmatrix} (v, Z) = e(-\frac{\pi}{2}B(v, v) + \pi i {}^t c^1 c^2) \theta_{Zw^1}^c(v)$
 - b) $\theta^{c+w}(v) = e(-\pi i {}^t w^1 c^2 + \pi i {}^t c^1 w^2 + \pi i {}^t w^1 w^2) \theta_w^c(v)$.
 - c) $\theta_{Zw^1 + w^2}^c(v) = \theta_{Zw^1}^c(v)$.
 - d) $\theta \begin{bmatrix} c^1 + w^1 \\ c^2 + w^2 \end{bmatrix} (v, Z) = e(2\pi i {}^t c^1 w^2 + 2\pi i {}^t w^1 w^2) \theta \begin{bmatrix} c^1 + w^1 \\ c^2 \end{bmatrix} (v, Z)$
6. The set $\langle \theta \begin{bmatrix} c^1 + D^{-1}m \\ c^2 \end{bmatrix}, m \in \mathbb{Z}_D \rangle$ is a base for classical theta functions for $L(H, \chi)$, where $c = Zc^1 + c^2$ is the characteristic of $L(H, \chi)$.

1.6 Action of the symplectic group.

Let D denote a polarization type. Let $\Gamma_D := \{R \in M_{2g}(\mathbb{Z}), R \begin{pmatrix} 0 & D \\ -D & 0 \end{pmatrix} {}^t R = \begin{pmatrix} 0 & D \\ -D & 0 \end{pmatrix}\}$.

Let $G_D := \{M \in \text{Sp}_{2g}(\mathbb{Q}), M = \begin{pmatrix} I & 0 \\ 0 & D \end{pmatrix}^{-1} R \begin{pmatrix} I & 0 \\ 0 & D \end{pmatrix}, \text{ for some } R \in \Gamma_D\}$. Note that for $R = \begin{pmatrix} A & B \\ \Gamma & \Delta \end{pmatrix} \in \Gamma_D$ and $M = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in G_D$ defined as above, we have

$$\alpha = A, \beta = BD, \gamma = D^{-1}\Gamma, \delta = D^{-1}\Delta D.$$

The group G_D is characterized as $G_D = \{M \in \text{Sp}_{2g}(\mathbb{Q}), {}^t M \Lambda_D \subset \Lambda_D\}$. The group G_D acts on $\mathfrak{h}_g$ as follows, see Ch.8, & 1 in [4]. Let $M = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in G_D$; we define $M(Z) := (\alpha Z + \beta D)(\gamma Z + \delta)^{-1}$. Two abelian varieties X_Z and $X_{Z'}$ of type D are isomorphic iff the matrices Z and Z' lie in the same orbit under the above action i.e. $Z' = M(Z)$. The isomorphism is given by $\phi(M) : X_Z \longrightarrow X_{M(Z)}$ s.t. the corresponding map $\phi(M)_r : \Lambda_Z \longrightarrow \Lambda_{M(Z)}$ has matrix $R = {}^t M^{-1}$ w.r.t. the canonical symplectic bases defined by Z and $M(Z)$.

It turns out that the map $\phi(M)_a : \mathbb{C}^g \longrightarrow \mathbb{C}^g$ has matrix A^{-1} , where $A = {}^t(\gamma Z + \delta)$, w.r.t. the standard base of $\mathbb{C}^g$. In addition, $\phi(M)_a^* H_{M(Z)} = H_Z$. We therefore have the following diagram

$$\begin{array}{ccccc}
\mathbb{R}^{2g} & \xrightarrow{j_Z} & \mathbb{C}^g & \longrightarrow & X_Z \\
\downarrow \phi(M)_r & & \downarrow \phi(M)_a & & \downarrow \phi(M) \\
\mathbb{R}^{2g} & \xrightarrow{j_{M(Z)}} & \mathbb{C}^g & \longrightarrow & X_{M(Z)}
\end{array} \tag{1}$$

Define $M_Z(v) := A^{-1}v (= \phi(M)_a(v))$. We then have that the action of G_D on $\mathfrak{h}_g$, lifts to an action on $\mathbb{C}^g \times \mathfrak{h}_g$ by defining $M(v, Z) := (M_Z(v), M(Z))$. Over the space $\mathfrak{h}_g$ there exists a universal family $\mathfrak{X}_D \longrightarrow \mathfrak{h}_g$ with fiber over Z to be the abelian variety X_Z . This is defined as follows. $\Lambda_D = \mathbb{Z}^g \oplus D\mathbb{Z}^g$ acts on $\mathbb{C}^g \times \mathfrak{h}_g$ by $n(v, Z) = (v + j_Z(n), Z)$. This action is free and properly discontinuous and the quotient space is $\mathfrak{X}_D$. The above action of G_D on $\mathbb{C}^g \times \mathfrak{h}_g$, descends to an action on $\mathfrak{X}_D$.

1.7 Factors of automorphy and line bundles - Sections and theta functions.

Let $\pi : V \longrightarrow X$ denote the canonical map. A factor of automorphy $f : \Lambda \times V \rightarrow \mathbb{C}^\times$ gives rise to a line bundle on X as follows: Take the action of Λ on $V \times \mathbb{C}$ defined by $\lambda(v, z) := (v + \lambda, f(\lambda, v)z)$. We then define L to be the quotient space of $V \times \mathbb{C}$ under this action.

We can also find transition functions for this line bundle: Take a covering $X = \cup_a U^a$ of X such that the preimage of each U^a under the map π to be a disjoint union of sets isomorphic to U^a . For each a , we choose one of those lifts, say W^a , and let $\pi_a : W^a \xrightarrow{\cong} U^a$ be the restriction of the map π . Define $w_a(x) := \pi_a^{-1}(x) \in W_a$. For any a, b with $U^{ab} := U^a \cap U^b \neq \emptyset$, there is a unique $\lambda_{ab} \in \Lambda$ satisfying $\lambda_{ab} := w_b(x) - w_a(x)$ for all $x \in U^{ab}$. In that way we get an identification of $W^a \cap \pi^{-1}(U^{ab})$ with $W^b \cap \pi^{-1}(U^{ab})$. Define now holomorphic functions $g_{ab} : U^{ab} \rightarrow \mathbb{C}^\times$ given by $g_{ab}(x) := f(\lambda_{ab}, w_a(x))$, which satisfy the cocycle condition for transition functions. Let $L^{tr} := \bigsqcup_a U^a \times \mathbb{C} / \sim$, where $(x_a, z_a) \sim (x_b, z_b)$ iff $x_a = x = x_b$ and $z_b = g_{ab}(x)z_a$, be the standard construction of a line bundle by its transition functions. There is an isomorphism of L^{tr} with L given by $L^{tr} \ni (x_a, z) \mapsto [w_a(x_a), z] \in L$ with inverse $L \ni [v, z] \mapsto (x_a = \pi(v), f(v - w_a(x_a), w_a(x_a))^{-1}z) \in L^{tr}$. In that way the g_{ab} 's become transition functions of L w.r.t. the trivialization induced by composing the natural trivializations of L^{tr} with the above defined inverse map. Note that by making different choices of the liftings W^a of U^a in the above, we get equivalent transition functions.

Let f and f' be equivalent factors of automorphy satisfying $f'(\lambda, v) = f(\lambda, v)h(v)h(v + \lambda)^{-1}$, where $h : V \rightarrow \mathbb{C}^\times$ holomorphic. By fixing the above covering data we define $h^a : U^a \rightarrow \mathbb{C}^\times$ by $h_a(x_a) := h(w_a(x_a))$. The line bundles L and L' corresponding to the above factors of automorphy are then isomorphic and their transition functions (constructed as above w.r.t. the common fixed covering data) satisfy $g'_{ab}(x) = g_{ab}(x) \frac{h_a(x)}{h_b(x)}$. There is also a canonical choice of isomorphism $\Phi_h : L \rightarrow L'$: By viewing the line bundles as quotient spaces, via the action of the lattice, the isomorphism is defined by $[v, z] \mapsto [v, h(v)^{-1}z]$. By viewing the line bundles as quotient spaces, via their transition functions and the above identification, the isomorphism is defined by sending $(x_a, z_a) \mapsto (x_a, h_a(x_a)^{-1}z_a)$.

Let $\phi : X' \rightarrow X$ be a map of abelian varieties. If f is a factor of automorphy for X , we denote by ϕ^*f the factor of automorphy for X' given by $(\phi_r \times \phi_a)^*f$. If L is the line bundle on X corresponding to f then ϕ^*L is the line bundle on X' corresponding to ϕ^*f . If θ is a theta function for the factor of automorphy f then $\phi_a^*\theta$ (or $\phi^*\theta$ in a simplified notation) is a theta function for the factor of automorphy ϕ^*f .

Sections of the line bundle L given by the factor of automorphy f , correspond to theta functions $\theta : V \rightarrow \mathbb{C}$ satisfying the functional equation $\theta(\lambda + v) = f(\lambda, v)\theta(v)$. The relation is the following. Given a section s of L , let $s(x) = [v_s(x), z_s(x)]$ (we view L as a quotient space given by the action of the lattice corresponding to f). We then define $\theta_s(v) := f(v - v_s(x), v_s(x))z_s(x)$. It turns out that this is a well defined theta function for the factor of automorphy f . Conversely, given a theta function θ for f , we define $s(x) := [v(x), \theta(v(x))]$, where $v(x)$ an arbitrary vector which lies over x . This is also well defined and the two processes are inverse to each other.

If $f' = f \star h$ as in Section 1.7 and $\Phi_h : L \rightarrow L'$ the canonical choice of isomorphism as in Section 1.7 above, then given a section $s' \in H^0(X, L')$ corresponding to the theta function $\theta_{s'}$, we have that $s := \Phi_h^*s' \in H^0(X, L)$ corresponds to the theta function $\theta_s := h(v)\theta_{f'}(v)$.

1.8 Theta transformation formulas.

Let X_Z be an abelian variety with a line bundle $L(H_Z, \chi)$ which defines a polarization H_Z on X_Z . We denote by c the characteristic of L . Let $M \in G_D$ as in Section 1.6 and define $Z' = M(Z)$. Let $\psi = \psi(M) : X_{Z'} \rightarrow X_Z$ be the inverse of the map $\phi = \phi(M) : X_Z \rightarrow X_{Z'}$ defined in Section 1.6. Then the line bundle $\psi^*L(H_Z, \chi)$ has type $\psi^*H_Z = H_{Z'}$, semicharacter $\chi' = \psi^*\chi$ and characteristic $M[c]$ with $M[c]^1 = \delta c^1 - \gamma c^2 + \frac{1}{2}(D\gamma^t \delta)_0$ and $M[c]^2 = -\beta c^1 + \alpha c^2 + \frac{1}{2}(\alpha^t \beta)_0$, see Lemma 4.1 of Ch. II in [4], where there is an unfortunate

omission of D in the expression of $M[c]^1$ (The $(\)_0$ stands for the diagonal vector of the matrix).

Lemma 1.1 *We have that $\psi^*e_{(H_Z, \chi)} = e_{(H_{Z'}, \chi')} \star h_\psi$, where $h_\psi(v) = e(\pi i {}^t v(\gamma Z + \delta)^{-1} \gamma v)$. Also, $\phi^*e_{(H_{Z'}, \chi')} = e_{(H_Z, \chi)} \star h$, where $h(v) = e(-\pi i {}^t v(\gamma Z + \delta)^{-1} \gamma v)$.*

Proof. We have that $a_{(H_Z, \chi)} = e_{(H_Z, \chi)} \star h_1$, where $h_1(v) = e(-\frac{\pi}{2} {}^t v(\text{Im} Z)^{-1} v)$, see Section 1.5. By applying ψ^* we get that $\psi^*a_{(H_Z, \chi)} = \psi^*e_{(H_Z, \chi)} \star \psi^*h_1$. Since $\psi^*a_{(H_Z, \chi)} = a_{(H_{Z'}, \chi')}$ and $a_{(H_{Z'}, \chi')} = e_{(H_{Z'}, \chi')} \star h'_1$, where $h'_1(v') = e(-\frac{\pi}{2} {}^t v'(\text{Im} Z)^{-1} v')$, we get that $h_\psi(v) = \psi^*h_1(v)^{-1} h'_1(v')$ i.e. $h_\psi(v) = e(\frac{\pi}{2} {}^t v(\text{Im} Z)^{-1} v - \frac{\pi}{2} {}^t v'(\text{Im} Z')^{-1} v')$, where $v' := \phi_a(v)$. A formula given in the proof of Proposition 6.1, Ch. 8 in [4] (or a straight forward calculation), implies the above form for h_ψ . To prove the second formula, we apply ϕ^* to the first one. $\square$

The tuple $B^Z := \langle \theta_{ZD^{-1}m}^c(v); m \in \mathbb{Z}_D \rangle$ forms a base of the canonical theta functions for $L(H_Z, \chi)$ and the tuple $B^{Z'} := \langle \theta_{Z'D^{-1}n}^{M[c]}(v'); n \in \mathbb{Z}_D \rangle$ forms a base of the canonical theta functions for $L(H_{Z'}, \chi')$. Since $\psi^*L(H_Z, \chi) = L(H_{Z'}, \chi')$ and $\psi^*a_{(H_Z, \chi)} = a_{(H_{Z'}, \chi')}$, we get that $\psi^*B^Z := \langle \psi_a^* \theta_{ZD^{-1}m}^c(v); m \in \mathbb{Z}_D \rangle$ forms also a base of the canonical theta functions for $L(H_{Z'}, \chi')$. There is then an invertible complex matrix C relating ψ^*B^Z with the base $B^{Z'}$.

Proposition 1.1 *Keeping the above notation, let we assume that the characteristic $c \in \frac{1}{2}\Lambda(H_Z)$. We then have that the matrix C with $\psi^*B^Z = C B^{Z'}$ is of the form $C = (\det(\gamma Z + \delta))^{-\frac{1}{2}} C_M$, where C_M is a matrix depending on M with $\det C_M = \zeta_8$, except when $3|d_g$ and $(d_{g-1}, 3) = 1$, in which case we have $\det C_M = \zeta_{24}$ (here by ζ_m we denote an m -root of unity).*

Proof. Let $G_D^{\text{int}} := G_D \cap \text{Sp}_{2g}(\mathbb{Z})$. A matrix M belongs to G_D^{int} if $M = \begin{pmatrix} I & 0 \\ 0 & D \end{pmatrix}^{-1} R \begin{pmatrix} I & 0 \\ 0 & D \end{pmatrix}$, where $R = \begin{pmatrix} A & B \\ \Gamma & \Delta \end{pmatrix} \in \Gamma_D$ with $\Gamma = D\Gamma_1$, $\Gamma_1 \in M_g(\mathbb{Z})$. Let therefore define $\Gamma_D^{\text{int}} := \{R = \begin{pmatrix} A & B \\ \Gamma & \Delta \end{pmatrix} \in \Gamma_D \text{ with } \Gamma = D\Gamma_1, \Gamma_1 \in M_g(\mathbb{Z})\}$. We have the following lemma:

Lemma 1.2 *The group Γ_D is generated by Γ_D^{int} and the matrix $J := \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix}$ and therefore the group G_D is generated by G_D^{int} and the matrix $\begin{pmatrix} 0 & -D \\ D^{-1} & 0 \end{pmatrix}$.*

Proof. For the proof we use results from [2]. We have seen that $K(D) = D^{-1}\mathbb{Z}_D \oplus \mathbb{Z}_D$. A matrix $R \in \Gamma_D$ defines an action on $K(D)$ given by multiplication by $\begin{pmatrix} I & 0 \\ 0 & D \end{pmatrix} {}^t R \begin{pmatrix} I & 0 \\ 0 & D \end{pmatrix}^{-1}$. By using the identification $K(D) \cong \mathbb{Z}_D \oplus \mathbb{Z}_D$ via the isomorphism $D^{-1}\mathbb{Z}_D \oplus \mathbb{Z}_D \longrightarrow \mathbb{Z}_D \oplus \mathbb{Z}_D$ given by the matrix $\begin{pmatrix} D & 0 \\ 0 & I \end{pmatrix}$, the action of a matrix $R \in \Gamma_D$ on $K(D) \cong \mathbb{Z}_D \oplus \mathbb{Z}_D$ is given by multiplication by $\bar{D} {}^t R \bar{D}^{-1}$, where $\bar{D} := \begin{pmatrix} D & 0 \\ 0 & D \end{pmatrix}$. One can define on $K(D)$ an alternating form e^D , see [4], [2], and the above action becomes a symplectic action with respect to that form. We denote the symplectic group (w.r.t. e^D) of $K(D) \cong \mathbb{Z}_D \oplus \mathbb{Z}_D$ by $\text{Sp}(D)$. We then have an exact sequence

$$0 \longrightarrow \Gamma_D(D) \longrightarrow \Gamma_D \xrightarrow{\pi} \text{Sp}(D) \longrightarrow 0,$$

where $\pi(R) := \bar{D} {}^t R \bar{D}^{-1}$ and $\Gamma_D(D) := \{R \in \Gamma_D \mid R = I + \bar{D}A, A \in M_{2g}(\mathbb{Z})\}$ (that the sequence is exact on the right it is proven in [2]). Note that $\Gamma_D(D) \subset \Gamma_D^{\text{int}}$. Therefore it is enough to show that every element of $\text{Sp}(D)$ has a lift to an element of Γ_D which is a product of the matrix J and elements of Γ_D^{int} .

Following the notation of [2], we have that $A \in L_D$ iff $\bar{D} {}^t A \bar{D}^{-1} \in \Gamma_D$, where L_D is defined in Section 2 of [2]. A matrix $A \in L_D$ acts on $K(D)$ by multiplication. In the proof of Theorem 2 of [2], it is shown that a matrix $A \in \text{Sp}(D)$ has a lift $\tilde{A} \in L_D$ which satisfies a relation of the form

$$\begin{pmatrix} I & 0 \\ c_1 & I \end{pmatrix} \begin{pmatrix} a & Dy \\ -D & d \end{pmatrix} \begin{pmatrix} U_1 & 0 \\ 0 & U_2 \end{pmatrix} \begin{pmatrix} I & b_1 \\ 0 & I \end{pmatrix} \tilde{A} = \begin{pmatrix} I & b_2 \\ 0 & I \end{pmatrix},$$

where y is an integral diagonal matrix and all the matrices belong to L_D . The inverse of a matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ in L_D belongs to L_D and is given by $\bar{D} \begin{pmatrix} {}^t d & -{}^t b \\ -{}^t c & {}^t a \end{pmatrix} \bar{D}^{-1}$. Therefore

A has a lift R in Γ_D which is given by $R = \bar{D} {}^t \tilde{A} \bar{D}^{-1}$. By observing now that $\begin{pmatrix} I & 0 \\ c & I \end{pmatrix} = J \begin{pmatrix} -I & c \\ 0 & -I \end{pmatrix} J$, we conclude that $R = (JN_1J)N_2N_3N_4(JN_5J)$, where $N_i \in \Gamma_D^{\text{int}}$. This proves the lemma. □

As in [6], we can rewrite the formula we want to prove in the form

$$\begin{aligned} &< \theta_{ZD^{-1}m}^c(v, Z); m \in \mathbb{Z}_D > \sqrt{dv_1 \wedge \dots \wedge dv_g} = \\ &= C_M < \theta_{Z'D^{-1}n}^{M[c]}(v', Z'); n \in \mathbb{Z}_D > \sqrt{dv'_1 \wedge \dots \wedge dv'_g}, \end{aligned}$$

where $v' = {}^t(\gamma Z + \delta)^{-1}v$ and $Z' = (\alpha Z + \beta)(\gamma Z + \delta)^{-1}$. Note that if $c \in \frac{1}{2}\Lambda(H_Z)$ then $M[c] \in \frac{1}{2}\Lambda(H_{Z'})$. We conclude that if the formula holds for $M_1, M_2 \in G_D$ then it also holds for $M_1 M_2$. It is enough therefore to verify the formula for matrices in the generators. We express the formula we want to prove in terms of classical theta functions by using Lemma 1.1 and the formulas in Section 1.5. It becomes

$$\begin{aligned} &< \theta \left[\begin{matrix} c^1 + D^{-1}m \\ c^2 \end{matrix} \right] (v, Z); m \in \mathbb{Z}_D > = e(-\pi i {}^t v (\gamma Z + \delta)^{-1} \gamma v) \\ &e(-\pi i {}^t M[c]^1 M[c]^2 + \pi i {}^t c^1 c^2) (\det(\gamma Z + \delta))^{-\frac{1}{2}} C_M < \theta \left[\begin{matrix} M[c]^1 + D^{-1}n \\ M[c]^2 \end{matrix} \right] (v', Z'); n \in \mathbb{Z}_D >. \end{aligned} \quad (2)$$

Matrices of the form $\begin{pmatrix} 0 & -D \\ D^{-1} & 0 \end{pmatrix}$. In this case $e(-\pi i {}^t v (\gamma Z + \delta)^{-1} \gamma v) = e(-\pi i {}^t v Z^{-1} v)$ and $\det(\gamma Z + \delta) = \frac{\det Z}{d}$. We also have that $v' = DZ^{-1}v$, $Z' = -DZ^{-1}D$ and $M[c]^1 = -D^{-1}c^2$, $M[c]^2 = Dc^1$. The relation (2) in this case becomes

$$\begin{aligned} &< \theta \left[\begin{matrix} c^1 + D^{-1}m \\ c^2 \end{matrix} \right] (v, Z); m \in \mathbb{Z}_D > = e(-\pi i {}^t v Z^{-1} v) \\ &e(2\pi i {}^t c^1 c^2) \left(\frac{\det Z}{d} \right)^{-\frac{1}{2}} C_M < \theta \left[\begin{matrix} -D^{-1}c^2 + D^{-1}n \\ Dc^1 \end{matrix} \right] (v', Z'); n \in \mathbb{Z}_D >. \end{aligned}$$

As in [6], we apply Fourier transform. Write $\theta \left[\begin{matrix} c^1 + D^{-1}m \\ c^2 \end{matrix} \right] (v, Z) = \sum_{\lambda \in \mathbb{Z}^g} f(\lambda)$ where $f(x) := e(\pi i {}^t (x + c^1 + D^{-1}m) Z (x + c^1 + D^{-1}m) + 2\pi i {}^t (v + c^2) (x + c^1 + D^{-1}m))$. Let $\hat{f}(x) := \int_{\mathbb{R}^g} f(x) e(2\pi i {}^t x \lambda) dx$. We then have that $\theta \left[\begin{matrix} c^1 + D^{-1}m \\ c^2 \end{matrix} \right] (v, Z) = \sum_{\lambda \in \mathbb{Z}^g} \hat{f}(\lambda)$. Using Lemma 5.8 of Ch. II in [6], we get by substituting $x' = (x + c^1 + D^{-1}m)$ that

$$\hat{f}(\lambda) = e(-2\pi i {}^t (c^1 + D^{-1}m) \lambda) \left(\det \frac{Z}{i} \right)^{-\frac{1}{2}} e(-\pi i {}^t (v + c^2 + \lambda) Z^{-1} (v + c^2 + \lambda)).$$

We therefore get that

$$\begin{aligned} &\theta \left[\begin{matrix} c^1 + D^{-1}m \\ c^2 \end{matrix} \right] (v, Z) = e(-\pi i {}^t (v + c^2) Z^{-1} (v + c^2)) \left(\det \frac{Z}{i} \right)^{-\frac{1}{2}} \\ &\sum_{\lambda \in \mathbb{Z}^g} e(-2\pi i {}^t (c^1 + D^{-1}m) \lambda - 2\pi i {}^t (v + c^2) Z^{-1} \lambda - \pi i {}^t \lambda Z^{-1} \lambda). \end{aligned}$$

Observe now that by substituting $-\lambda = Dk + n$ with $n \in \mathbb{Z}_D$, we can rewrite the sum as

$$\begin{aligned} &\sum_{\lambda \in \mathbb{Z}^g} e(-2\pi i {}^t (c^1 + D^{-1}m) \lambda - 2\pi i {}^t (v + c^2) Z^{-1} \lambda - \pi i {}^t \lambda Z^{-1} \lambda) = \\ &\sum_{n \in \mathbb{Z}_D} \sum_{k \in \mathbb{Z}^g} e \left(2\pi i {}^t (c^1 + D^{-1}m) (Dk + n) + 2\pi i {}^t (v + c^2) Z^{-1} (Dk + n) - \pi i {}^t (Dk + n) Z^{-1} (Dk + n) \right). \end{aligned}$$

Using straight forward calculation, one finally deduces that

$$\theta \left[\begin{array}{c} c^1 + D^{-1}m \\ c^2 \end{array} \right] (v, Z) = e(-\pi i {}^t v Z^{-1} v) \left(\det \frac{Z}{i} \right)^{-\frac{1}{2}} e(2\pi i {}^t c^1 c^2) \\ \sum_{n \in \mathbb{Z}_D} e(2\pi i {}^t m D^{-1} n) \theta \left[\begin{array}{c} -D^{-1}c^2 + D^{-1}n \\ Dc^1 \end{array} \right] (v', Z').$$

The matrix C_M we are asking for has m, n entry equal to $(\frac{d}{ig})^{-\frac{1}{2}} e(2\pi i {}^t m D^{-1} n)$. Let $d := \det D$. The matrix $A = (a_{mn})_{m, n \in \mathbb{Z}_D}$ with $a_{mn} := e(2\pi i {}^t m D^{-1} n)$ has determinant $\det A = \zeta_4 d^{\frac{d}{2}}$. One way to see this is the following. Denote by $\mathbb{C}[\mathbb{Z}_{d_i}]$ the $\mathbb{C}$ -vector space of dimension d_i “corresponding” to the finite group $\mathbb{Z}_{d_i}$. Fix the natural base $< m, m \in \mathbb{Z}_{d_i} >$ on that vector space. Define the map $\phi_i : \mathbb{C}[\mathbb{Z}_{d_i}] \rightarrow \mathbb{C}[\mathbb{Z}_{d_i}]$ by $\phi_i(m) := \sum_{n \in \mathbb{Z}_{d_i}} e(2\pi i n d_i^{-1} m) n$. Let C_i be the matrix corresponding to ϕ_i . Then $\det C_i = \zeta_4 d_i^{\frac{d_i}{2}}$, where ζ_4 is 4-th root of unity. Indeed, it is easy to see that $\det(C_i^2) = \pm d_i^{d_i}$. Observe now that the matrix A is the matrix corresponding to the tensor product of the maps ϕ_i and so, its determinant is $\det A = \det C_1^{\frac{d}{d_1}} \cdots \det C_g^{\frac{d}{d_g}} = \zeta_4 d^{\frac{d}{2}}$.

To conclude this case, observe that $(\frac{d}{ig})^{-\frac{1}{2}} = \zeta_8 d^{-\frac{d}{2}}$. Therefore $\det C_M = \zeta_8$.

Matrices in G_D^{int} . Let $M \in G_D^{\text{int}}$. Then M corresponds to an isomorphism $\psi : X_{Z'}/\Lambda_{Z'} \rightarrow X_Z/\Lambda_Z$ which is a lift of an isomorphism of ppav. In this case the usual theta transformation formula holds, see Ch. 8, & 6 in [4]. Let $a = c + Zw^1$, $c \in \frac{1}{2}\Lambda(H)$, $w^1 = D^{-1}w_1 \in D^{-1}\mathbb{Z}_D$.

We have the following, see Ch. 8, & 4 and & 6 in [4].

1. $\psi^* \theta^a(v, Z) = C(Z, M, a) \theta^{M[a]}(v', Z')$.
2. $C(Z, M, a) = C(Z, M, 0) e(\pi i E(M[0], A^{-1}a))$.
3. $C(Z, M, 0) = k(M) e(\pi i {}^t M[0]^1 M[0]^2) \det(\gamma Z + \delta)^{-\frac{1}{2}}$, where $k(M)$ is an 8-th root of unity.

Note that $M[a]^1 = M[c]^1 + \delta w^1$ and $M[a]^2 = M[c]^2 - \beta w^1$. Using the above formulas and the formulas in Section 1.5 we get

$$e(-\pi i {}^t c^2 w^1) \psi^* \theta_{Zw^1}^c(v, Z) = C(Z, M, 0) e(\pi i E(M[0], A^{-1}a)) \\ e(-\pi i {}^t (\delta w^1) M[c]^2 + \pi i {}^t M[c]^1 (-\beta w^1) + \pi i {}^t (\delta w^1) (-\beta w^1)) \theta_{Z' \delta w^1}^{M[c]}(v', Z').$$

We have

$$M[0]^1 = \frac{1}{2}(D\gamma {}^t \delta)_0, \quad M[0]^2 = \frac{1}{2}(\alpha {}^t \beta)_0$$

$$\begin{aligned} (A^{-1}a)^1 &= \delta(c^1 + w^1) - \gamma c^2, & (A^{-1}a)^2 &= -\beta(c^1 + w^1) + ac^2 \\ M[c]^1 &= \delta c^1 - \gamma c^2 + \frac{D}{2}(\gamma^t \delta)_0, & M[c]^2 &= -\beta c^1 + ac^2 + \frac{1}{2}(\alpha^t \beta)_0. \end{aligned}$$

We then get

$$\psi^* \theta_{ZD^{-1}w_1}^c(v, Z) = k(M)e(\pi i k)(\pi i \lambda w^1)e(-\pi i^t w^1 \beta \delta w^1) \det(\gamma Z + \delta)^{-\frac{1}{2}} \theta_{Z'D^{-1}\Delta w_1}^{M[c]},$$

where $k = {}^t M[0]^1 M[0]^2 + {}^t M[0]^1(-\beta c^1 + ac^2) - {}^t M[0]^2(\delta c^1 - \gamma c^2)$ and $\lambda = -{}^t M[0]^1 \beta - {}^t M[0]^2 \delta - {}^t M[c]^2 \delta - {}^t M[c]^1 \beta + {}^t c^2$. Observe now that $k \in \frac{1}{4d_g} \mathbb{Z}^g$ and $\lambda \in \frac{1}{2} \mathbb{Z}^g$.

Note that when $\gamma \in M_g(\mathbb{Z})$ i.e. $\Gamma = D\Gamma_1$ for some integral matrix Γ_1 , we have that Δ acts as a permutation on $\mathbb{Z}_D$. Indeed, the relation $\Delta D^t A - \Gamma D^t B = D$ implies $\Delta(D^t A D^{-1}) = I + \Gamma(D^t B D^{-1})$ i.e. $\Delta(D^t A D^{-1}) = I + D\Gamma_1(D^t B D^{-1})$ and so, $\Delta A_1 = I + DM$ for some integral matrices A_1, M . This implies that Δ induces an epimorphism on the group $\mathbb{Z}_D$ and therefore an automorphism.

Therefore the matrix C_M of the proposition is going to have in the $w_1, \Delta w_1$ -entry the value $k(M)e(\pi i k)e(\pi i \lambda D^{-1} w_1)e(-\pi i^t w_1^t \Delta D^{-1} B w_1)$ and any other entry zero. We have

$$\prod_{w_1 \in \mathbb{Z}_D} e(\pi i \lambda D^{-1} w_1) = e\left(\pi i \sum_{i=1}^g \frac{\lambda_i}{d_i} \sum_{w_1 \in \mathbb{Z}_D} w_1^i\right) = e\left(\pi i \sum_{i=1}^g \frac{\lambda_i}{d_i} \frac{d}{d_i} \frac{d_i(d_i - 1)}{2}\right)$$

The sum which appears above belongs to $\frac{1}{4} \mathbb{Z}$ and so, the product is a ζ_8 . Also, the matrix ${}^t \Delta D^{-1} B$ is symmetric and let $\alpha_{ij} = \frac{a_{ij}}{d_i}, a_{ij} \in \mathbb{Z}$ its ij -entry. Then

$$\prod_{w_1 \in \mathbb{Z}_D} e(\pi i^t w_1^t \Delta D^{-1} B c_1) = e\left(\pi i \sum_{i=1}^g \frac{a_{ii}}{d_i} d \frac{(d_i - 1)(2d_i - 1)}{6} + 2 \sum_{1 \leq i < j \leq g} \frac{a_{ij}}{d_i} d \frac{(d_i - 1)(d_j - 1)}{4}\right)$$

The sums which appear above belong to $\frac{1}{2} \mathbb{Z}$ and so, the product is a ζ_4 , except when $3|d_g$ and $(d_{g-1}, 3) = 1$, in which case it belongs to $\frac{1}{6} \mathbb{Z}$ and the product is a ζ_{12} . To conclude, we have that $\det C_M = \zeta_8$, except when $3|d_g$ and $(d_{g-1}, 3) = 1$, in which case $\det C = \zeta_{24}$

□

For the case now of a totally symmetric bundle, note first that in Lemma 1.1, if $L(H_Z, \chi_0)$ is of characteristic $c = 0$, then $\psi^* L(H_Z, \chi_0)$ is again of characteristic 0. Indeed, in the case of an “even” polarization, we have always that $\chi_0 = 1$ and so, $\psi_r^* \chi_0 = 1 = \chi_0'$.

Proposition 1.2 *Keeping the notation of Proposition 1.1, we assume in addition that the characteristic $c = 0$, $\mathcal{L}$ is totally symmetric and that $g \geq 3$. We then have that the matrix*

C with $\psi^* B^Z = C B^{Z'}$ is of the form $C = (\det(\gamma Z + \delta))^{-\frac{1}{2}} C_M$, where C_M is a matrix depending on M with $\det C_M = 1$, except when $3|d_g$ and $(d_{g-1}, 3) = 1$ in which case we have $\det C_M = \zeta_3$.

Proof. The proof is a modification of the proof of Proposition 1.1.

At the end of the paragraph “**Matrices of the form** $\begin{pmatrix} 0 & -D \\ D^{-1} & 0 \end{pmatrix}$ ”: For $g \geq 3$ the number $\frac{d}{d_i}$ is a multiple of 4 and so, $\det C_i^{\frac{d}{d_i}} = d_i^{\frac{d_i}{2}}$. Therefore $\det A = d^{\frac{d}{2}}$. Also, for $g \geq 3$ we have $(\frac{d}{d_g})^{-\frac{d}{2}} = d^{-\frac{d}{2}}$. Therefore, $\det C_M = 1$.

At the end of the paragraph “**Matrices in G_D^{int} ”**”: The first sum is an even integer and so, the first product is 1. For the second sum, if $g \geq 3$ the right summand is an even integer. For the left summand we have that it is an even integer except when $3|d_g$ and $(d_{g-1}, 3) = 1$ in which case it belongs to $\frac{2}{3}\mathbb{Z}$. Therefore for the second product we have that it is 1, except when $3|d_g$ and $(d_{g-1}, 3) = 1$, in which case it is ζ_3 . We also claim that $\text{sgn}(\Delta) = 1$, where by $\text{sgn}(\Delta)$ we denote the sign of the permutation of $\mathbb{Z}_D$ defined by the action of Δ .

Indeed, let $d_i = 2^{k_i} m_i$ with $1 \leq k_1 \leq k_2 \leq \dots \leq k_g$ and $m_1|m_2|\dots|m_g$ odd integers. Define $\mathbb{Z}_{\text{ev}} := \mathbb{Z}_{2^{k_1}} \oplus \dots \oplus \mathbb{Z}_{2^{k_g}}$ a group of order $n_{\text{ev}} = 2^{k_1+\dots+k_g}$ and $\mathbb{Z}_{\text{odd}} := \mathbb{Z}_{m_1} \oplus \dots \oplus \mathbb{Z}_{m_g}$ a group of order $n_{\text{odd}} = m_1 \dots m_g$. Then $\mathbb{Z}_D = \mathbb{Z}_{\text{ev}} \oplus \mathbb{Z}_{\text{odd}}$. Let $\phi : \mathbb{Z}_D \longrightarrow \mathbb{Z}_D$ an automorphism. Then $\phi(\mathbb{Z}_{\text{ev}}) = \mathbb{Z}_{\text{ev}}$ and $\phi(\mathbb{Z}_{\text{odd}}) = \mathbb{Z}_{\text{odd}}$. Indeed, $\mathbb{Z}_{\text{ev}} = \{x \in \mathbb{Z}_D \text{ with } 2^{k_g}x = 0\}$ and $\mathbb{Z}_{\text{odd}} = \{x \in \mathbb{Z}_D \text{ with } m_g x = 0\}$. Both those relations are invariant under ϕ . Let therefore ϕ_{ev} be the restriction of ϕ to $\mathbb{Z}_{\text{ev}}$ and ϕ_{odd} the restriction of ϕ to $\mathbb{Z}_{\text{odd}}$. If we interpret ϕ as a linear automorphism of the vector space $\mathbb{C}[\mathbb{Z}_D]$ then $\phi = \phi_{\text{ev}} \otimes \phi_{\text{odd}}$. By using the formula for the determinant of the tensor product of operators, we get $\text{sgn}\phi = \text{sgn}\phi_{\text{ev}}^{n_{\text{odd}}} \text{sgn}\phi_{\text{odd}}^{n_{\text{ev}}}$. Therefore, since n_{ev} is an even number, it is enough to prove the result for $\mathbb{Z}_D = \mathbb{Z}_{\text{ev}}$.

Let now E be the matrix which corresponds to the automorphism ϕ_{ev} . We call elementary transformations of $\mathbb{Z}_{\text{ev}}$ those which correspond to left or right multiplication by a matrix of one of the following forms: 1 in the diagonal and $a_{ij} \in \mathbb{Z}$ in some ij -entry if $j \geq i$ (and zero every where else) or 1 in the diagonal and $2^{k_i-k_j}a_{ij}$, $a_{ij} \in \mathbb{Z}$, in some ij -entry if $j > i$ (and zero every where else). Note that both those types of matrices induce automorphisms of $\mathbb{Z}_{\text{ev}}$. We then claim that by multiplying the matrix $E = (e_{ij})$ with the above type of matrices we can derive a matrix with all the elements of the last row, except the diagonal one, to be zero mod 2^{k_g} and the i -th element of the last column, with $1 \leq i < g$, to be zero mod 2^{k_i} . Indeed, we can first assume that e_{gg} is an odd integer: the determinant of E must be an odd number in order E to define an automorphism and so, some of the elements of the last row must

be odd. If e_{gg} is even, let e_{gj_0} , $j_0 < g$ be the odd element. But then using an elementary transformation we can add the j_0 -th column to the last column and so the gg -entry becomes odd. Since $D^{-1}ED$ is an integral matrix we have that $e_{gj} = 2^{k_g - k_j} m_{gj}$, $m_{gj} \in \mathbb{Z}$. But now the equation $2^{k_g - k_j} e_{gg} x \equiv -2^{k_g - k_j} m_{gj} \pmod{2^{k_g}}$ has a solution and therefore by multiplying the matrix E on the right by the elementary matrix with $2^{k_g - k_j} x$ in the gj -entry, we get that the gj -entry of the product is zero mod 2^{k_g} . By a similar argument, by multiplying on the left by an elementary matrix with x in the ig -entry, where x is the solution of $c_{gg} x \equiv -c_{ig} \pmod{2^{k_i}}$, we get that the ig -entry of the product is zero mod 2^{k_i} .

A matrix like the one we produced, corresponds to an even permutation of $\mathbb{Z}_{\text{ev}}$. Indeed, by writing $\mathbb{Z}_{\text{ev}}$ as a direct sum of two groups, the second of which is the $\mathbb{Z}_{2^{k_g}}$, the action is a direct sum of actions. Then, by applying the formula for the signatures stated above, we get that the signature of the permutation is one, since both groups are of even order. A similar argument gives that the action which is given by the elementary matrices induces an even permutation (here we have to use the hypothesis that $g \geq 3$). We therefore get that the permutation given by ϕ_{ev} is an even one. This concludes the proof of the proposition. $\square$

2 Abelian fibrations.

Everything we have stated which holds for a fixed abelian variety $X = V/\Lambda$, can be transferred easily to hold for a fibration $X \longrightarrow U$ of abelian varieties of type D , with the base U to be a simply connected Stein manifold (such as $\mathfrak{h}_g$). In this case, the universal covering $\tilde{X}$ of X will take the place of V and the homotopy group $\pi_1(X)$ will take the place of Λ . For the universal abelian fibration $f_{\text{un}} : \mathfrak{X}_D \longrightarrow \mathfrak{h}_g$ we have by construction, see Section 1.6, that $\pi_1(\mathfrak{X}_D) = \Lambda_D$ and $\tilde{\mathfrak{X}}_D = \mathbb{C}^{2g} \times \mathfrak{h}_g$. On $\mathfrak{X}_D$ we have a line bundle corresponding to the (holomorphic) classical factor of automorphy $e : \Lambda_D \times (\mathbb{C}^g \times \mathfrak{h}_g) \longrightarrow \mathbb{C}^\times$ given by $e(l; v, Z) = e(-\pi i^t \lambda^1 Z \lambda^1 - 2\pi i^t v \lambda^1)$, where λ^1 denotes the first g components of l . We denote by $\mathcal{L}_{\mathfrak{X}}$ the above line bundle.

2.1 Canonical form of line bundles on abelian fibrations.

Let $f : X \longrightarrow B$ be a fibration of abelian varieties with a symmetric line bundle $\mathcal{L}$ on X , trivialized along the zero section, which defines a polarization of type D on each fiber. We denote by $s : B \longrightarrow X$ its zero section and let $S := s(B)$. We cover S by open analytic (connected) sets V^a of X . The sets $U^a := s^* V^a$ define an open covering of B . Let

$(U^a, \lambda_1^a, \dots, \lambda_{2g}^a)$ be a choice of a symplectic base on the fibers of the restriction of the abelian fibration over U^a . For each a, b with $U^{ab} \neq \emptyset$, there is a matrix $M^{ab} = \begin{pmatrix} \alpha^{ab} & \beta^{ab} \\ \gamma^{ab} & \delta^{ab} \end{pmatrix} \in G_D$ relating the two symplectic bases as in Section 1.6. Let X^a be the preimage of U^a under f . We then have a period map $p_a : U^a \rightarrow \mathfrak{h}_g$ such that the diagram

$$\begin{array}{ccc} \tilde{X}^a & \xrightarrow{\tilde{\pi}_a} & \mathbb{C}^g \times \mathfrak{h}_g \\ \downarrow & & \downarrow \\ X^a & \xrightarrow{\pi_a} & \mathfrak{X}_D \\ \downarrow f & & \downarrow f_{\text{un}} \\ U^a & \xrightarrow{p_a} & \mathfrak{h}_g \end{array} \quad (3)$$

is a diagram of fiber products, where $\tilde{X}^a$ denotes the universal cover of X^a . We put $Z^a(s) := p_a(s)$, $s \in U^a$. Having chosen the period map p_a , we can identify $\tilde{X}^a$ with $\mathbb{C}^g \times p_a(U^a)$ and $\pi_1(X^a)$ with Λ_D . We write $(\pi_1(X^a), \tilde{X}^a) \cong^{p_a} (\Lambda_D, \mathbb{C}^g \times p_a(U^a))$ to denote this identification. We now “complete” V^a to an open cover $X^a = \cup_i V_i^a$ of X^a s.t. $V_i^a \cap S = \emptyset$ for all $V_i^a \neq V^a$ and such that each V_i^a has an isomorphic lift W_i^a in the universal cover $\tilde{X}^a \cong^{p_a} \mathbb{C}^g \times p_a(U^a)$. We can also assume that the intersection of any two of those open sets is simply connected.

Consider the induced diagram

$$\begin{array}{ccccccc} X^{ab} & \xrightarrow{\pi_a} & \pi_a(X^{ab}) & \xrightarrow{\phi_{ab}} & \pi_b(X^{ab}) & \xleftarrow{\pi_b} & X^{ab} \\ \downarrow f & & \downarrow f_{\text{un}} & & \downarrow f_{\text{un}} & & \downarrow f \\ U^{ab} & \xrightarrow{p_a} & p_a(U^{ab}) & \xrightarrow{\mu_{ab}} & p_b(U^{ab}) & \xleftarrow{p_b} & U^{ab} \end{array} \quad (4)$$

We denote by μ_{ab} the map which sends $Z \mapsto M^{ab}(Z)$ and by ϕ_{ab} the map induced by the action of M^{ab} on the universal abelian variety $\mathfrak{X}$ (fiberwise is the map $\phi(M)$ of Section 1.6). We have that $\pi_b = \phi_{ab} \pi_a$ and $p_b = \mu_{ab} p_a$.

Note that a symmetric line bundle $L(H, \chi)$ has always characteristic $c \in \frac{1}{2}\Lambda(H)$ w.r.t. any decomposition of H . Indeed, that $L(H, \chi)$ is symmetric is equivalent to $\chi(\lambda) = \pm 1$ for all $\lambda \in \Lambda$. The claim is a consequence of the definition of $\chi(\lambda)$. The restriction of $\mathcal{L}$ on the fibers over U^a is then of characteristic $c_a = Z^a(s)c_a^1 + c_a^2$, where $(c_a^1, c_a^2) \in \frac{1}{2}(D^{-1}\mathbb{Z}^g \oplus \mathbb{Z}^g)$, w.r.t. the decomposition defined by the choice of the symplectic base. Note that when $\mathcal{L}$ is totally symmetric the characteristic is going to be 0.

Define the line bundles $\mathcal{L}_a := \pi_a^* T_{c_a}^* \mathcal{L}_{\mathfrak{X}}$ on X^a . Note that in the totally symmetric case, we have that $\mathcal{L}_a := \pi_a^* \mathcal{L}_{\mathfrak{X}}$. Then $\mathcal{L}_a \cong \mathcal{L}|_{X^a}$ because the restriction on the fibers have the same type and characteristic and U^a is an open analytic set. Via the identification $(\pi_1(X^a), \tilde{X}^a) \xrightarrow{p_a} (\Lambda_D, \mathbb{C}^g \times p_a(U^a))$, the line bundle $\mathcal{L}_a$ corresponds to the factor of automorphy $e_a : \Lambda_D \times (\mathbb{C}^g \times U^a) \longrightarrow \mathbb{C}^\times$, where $e_a = e_a(\lambda^a, v^a, Z^a(s))$ is the classical factor of automorphy of characteristic c_a .

On X^{ab} we have $\mathcal{L}|_{X^{ab}} \cong \pi_b^* T_{c_b}^* \mathcal{L}_{\mathfrak{X}} \cong \pi_a^* \phi_{ab}^* (T_{c_b}^* \mathcal{L}_{\mathfrak{X}}|_{\pi_b(X^{ab})})$. On the other hand, $\mathcal{L}|_{X^{ab}} \cong \pi_a^* T_{c_a}^* \mathcal{L}_{\mathfrak{X}}$. We therefore have that $\phi_{ab}^* (T_{c_b}^* \mathcal{L}_{\mathfrak{X}}|_{\pi_b(X^{ab})}) \cong T_{c_a}^* \mathcal{L}_{\mathfrak{X}}|_{\pi_a(X^{ab})}$. Note that when we fix the identification $(\pi_1(X^{ab}), \tilde{X}^{ab}) \xrightarrow{p_a} (\Lambda_D, \mathbb{C}^g \times p_a(U^{ab}))$, the line bundle $\mathcal{L}_a|_{X^{ab}}$ corresponds to the factor of automorphy e_a while the line bundle $\mathcal{L}_b|_{X^{ab}}$ corresponds to the factor of automorphy $\phi_{ab}^* e_b$.

By Lemma 1.1, we have that $\phi_{ab}^* e_b|_{\Lambda_D \times \mathbb{C}^g \times p_a(U^{ab})} = e_a|_{\Lambda_D \times \mathbb{C}^g \times p_a(U^{ab})} \star h^{ab}$. In other words we have that

$$\phi_{ab}^* e_b = e_a \star h^{ab} \text{ on } (\pi_1(X^{ab}), \tilde{X}^{ab}) \xrightarrow{p_a} (\Lambda_D, \mathbb{C}^g \times p_a(U^{ab})),$$

where $h^{ab}(v^a, Z^a(s)) = e(-\pi i {}^t v^a (\gamma^{ab} Z^a(s) + \delta^{ab})^{-1} \gamma^{ab} v^a)$. Note that $v^b = {}^t (\gamma Z^a(s) + \delta)^{-1} v^a$ and $Z^b(s) = (\alpha Z^a(s) + \beta)(\gamma Z^a(s) + \delta)^{-1}$.

We glue now all the line bundles $(\mathcal{L}_a, X^a)$ together to get on X a line bundle $\mathcal{L}_{\text{can}}$ which restricts to the trivial bundle on the zero section of the fibration, in the following way:

Fix a trivialization $g_{ij}^a = e_a(\lambda_{ij}^a, w_i^a, Z^a(s))$ of $\mathcal{L}_a$ on X^a (resp. $g_{kl}^b = e_b(\lambda_{kl}^b, w_k^b, Z^b(s))$ of $\mathcal{L}_b$ on X^b), by choosing lifts W_i^a of the open sets V_i^a in the universal cover $\tilde{X}^a \xrightarrow{p_a} \mathbb{C}^g \times p_a(U^a)$ (resp. lifts W_k^b of V_k^b in the universal cover $\tilde{X}^b \xrightarrow{p_b} \mathbb{C}^g \times p_b(U^b)$). We then have a canonical isomorphism $\mathcal{L}_a \xrightarrow{\sim} \bigsqcup_i (V_i^a \times \mathbb{C}) / \sim \xrightarrow{\sim} \bigsqcup_i (V_i^a \times \mathbb{C}) / \sim$ (resp. $\mathcal{L}_b \xrightarrow{\sim} \bigsqcup_k (V_k^b \times \mathbb{C}) / \sim \xrightarrow{\sim} \bigsqcup_k (V_k^b \times \mathbb{C}) / \sim$), see Section 1.7.

Let now $\Phi_{ab} : \mathcal{L}_a|_{X^{ab}} \longrightarrow \mathcal{L}_b|_{X^{ab}}$ be the canonical isomorphism defined in Section 1.7 corresponding to $\phi_{ab}^* e_b = e_a \star h^{ab}$. We construct an explicit isomorphism ψ^{ab} which makes the following diagram commutative:

$$\begin{array}{ccc} \mathcal{L}_a|_{X^{ab}} & \xrightarrow{\Phi_{ab}} & \mathcal{L}_b|_{X^{ab}} \\ \downarrow \simeq & & \downarrow \simeq \\ \bigsqcup_i (V_i^a \times \mathbb{C}) / \sim|_{X^{ab}} & \xrightarrow{\psi^{ab}} & \bigsqcup_k (V_k^b \times \mathbb{C}) / \sim|_{X^{ab}} \end{array} \quad (5)$$

Such an isomorphism is given by $(x_i^a(s) = x, z_i^a) \mapsto (x_k^b(s) = x, z_k^b = \psi_{ik}^{ab}(x) z_i^a)$, where the ψ_{ik}^{ab} 's satisfy the relation $\psi_{jl}^{ab}(x) e_a(\lambda_{ij}^a, w_i^a(x), Z^a(s)) = e_b(\lambda_{kl}^b, w_k^b(x), Z^b(s)) \psi_{ik}^{ab}(x)$ and the cocycle conditions $\psi_{ik}^{ab} \psi_{ki}^{ba} = 1$, $\psi_{ik}^{ab} \psi_{ks}^{bc} \psi_{si}^{ca} = 1$ (we follow here the notation of Section 1.7).

Let $\pi_i^a : W_i^a \xrightarrow{\sim} V_i^a$ denote the canonical maps. For each $V_{ik}^{ab} := V_i^a \cap V_k^b \neq \emptyset$, we have that $\pi_i^{a-1}(V_{ik}^{ab}) \subset W_i^a$ is a lift of V_{ik}^{ab} in the universal cover $\tilde{X}^{ab} \xrightarrow{p_a} \mathbb{C}^g \times p_a(U^{ab})$. We also have that $\tilde{\phi}_{ab}^{-1} \pi_k^{b-1}(V_{ik}^{ab})$ is a lift of V_{ik}^{ab} in the same universal cover, where $\tilde{\phi}_{ab}$ is the map induced by ϕ_{ab} on the level of universal coverings (i.e. is the analytic representation of ϕ_{ab}). There exists therefore an element μ_{ik}^{ab} of Λ_D such that $\tilde{\phi}_{ab}^{-1} w_k^b(x) = \mu_{ik}^{ab} + w_i^a(x)$ for all $x \in V_{ik}^{ab}$.

We now define

$$\psi_{ik}^{ab}(x) = e_a(\mu_{ik}^{ab}, w_i^a(x), Z^a(s)) h^{ab} \left(\tilde{\phi}_{ab}^{-1}(w_k^b(x)), Z^a(s) \right)^{-1}.$$

It is a straight forward calculation to verify that those make the above diagram to commute and satisfy the required relations for automorphisms.

We define $\mathcal{L}_{\text{can}} := \bigsqcup_{a,i} (V_i^a \times \mathbb{C}) / \sim$, where $(x_i^a, z_i^a) \sim (x_k^b, z_k^b)$ iff

$$x_i^a = x_k^b = s \quad \text{and} \quad \begin{cases} z_k^b = g_{ik}^a z_i^a & \text{if } a = b \\ z_k^b = \psi_{ik}^{ab} z_i^a & \text{if } a \neq b \end{cases}.$$

Note that those satisfy the cocycle relation for transition functions. We also have that the restriction of $\mathcal{L}_{\text{can}}$ to the zero section S is trivial. Indeed, for $x \in V^{ab} \cap S$, we have that $w^a := w^a(x)$, $w^b := w^b(x) \in \Lambda_D$ and $\tilde{w}^b := \tilde{\phi}_{ab}^{-1}(w^b(x)) \in \Lambda_D$. Using the defining properties of the factors of automorphy and the definition of h^{ab} , we get that $e_a(\mu^{ab}, w^a(x), Z^a(s)) = e_a(\tilde{w}^b, 0, Z^a(s)) e_a(w^a, 0, Z^a(s))^{-1}$ and $h^{ab}(\tilde{w}^b, Z^a(s))^{-1} = e_b(w^b, 0, Z^b(s)) e_a(\tilde{w}^b, 0, Z^a(s))^{-1}$. We conclude that for $s \in S$, we have $\psi^{ab}(s) = \frac{f^a(s)}{f^b(s)}$, where $f^a : U^a \rightarrow \mathbb{C}^\times$ is the holomorphic function given by $f^a(s) = e(w^a, 0, Z^a(s))^{-1}$.

Finally observe that $\mathcal{L} \cong \mathcal{L}_{\text{can}}$. Indeed, they both restrict to isomorphic line bundles on the fibers (since they are both of type D and have the same characteristic) and they are both trivial along the zero section. The see-saw principle implies that they are isomorphic.

2.2 Proof of Theorems A and B.

It suffices to prove the theorem for the line bundle $\mathcal{L} := \mathcal{L}_{\text{can}}$ constructed in the previous section. The functions $\theta \left[\begin{smallmatrix} c_a^1 + D^{-1}n \\ c_a^2 \end{smallmatrix} \right] (v^a, Z^a(s))$, $n \in \mathbb{Z}_D$, are theta functions for the classical factor of automorphy $e_a : \Lambda_D \times \mathbb{C}^g \times \mathfrak{h}_g \rightarrow \mathbb{C}^\times$ of characteristic c_a . The line bundle $T_{c_a}^* \mathcal{L}_x|_{\pi_a(X^a)}$ corresponds by construction to the classical factor of automorphy e_a . Let $\tilde{\sigma}_n^a$ denote the section of $T_{c_a}^* \mathcal{L}_x|_{\pi_a(X^a)}$ corresponding to the theta function $\theta \left[\begin{smallmatrix} c_a^1 + D^{-1}n \\ c_a^2 \end{smallmatrix} \right] (v^a, Z^a(s))$. Then $\sigma_n^a := \pi_a^* \tilde{\sigma}_n^a$ is the section of $\mathcal{L}_a$ corresponding to the same theta function by identifying $(\pi_1(X^a), \tilde{X}^a) \xrightarrow{p_a} (\Lambda_D, \mathbb{C}^g \times p_a(U^a))$.

In the following, fix the identification $(\pi_1(X^{ab}), \tilde{X}^{ab}) \xrightarrow{p_a} (\Lambda_D, \mathbb{C}^g \times p_a(U^{ab}))$. The section $\phi_{ab}^*(\tilde{\sigma}_m^b|_{\pi_b(X^{ab})})$ is a section of the line bundle $\phi_{ab}^*(T_{c_b}^*\mathcal{L}_{\mathfrak{X}}|_{\pi_b(X^{ab})})$ corresponding to the theta function $\phi_{ab}^*\theta \left[\begin{smallmatrix} c_b^1 + D^{-1}m \\ c_b^2 \end{smallmatrix} \right] (v^b, Z^b(s))$ w.r.t. the factor of automorphy $\phi_{ab}^*e_b$. We therefore get that the section $\pi_a^*(\phi_{ab}^*\tilde{\sigma}_m^b|_{\pi_b(X^{ab})}) = \pi_b^*(\tilde{\sigma}_m^b|_{\pi_b(X^{ab})}) = \sigma_m^b|_{X^{ab}}$ is a section of $\mathcal{L}_b|_{X^{ab}}$ corresponding to the same theta function. Let $\Phi_{ab} : \mathcal{L}_a|_{X^{ab}} \longrightarrow \mathcal{L}_b|_{X^{ab}}$ be the canonical isomorphism defined in Section 2.1. Then $\Phi_{ab}^*\sigma_m^b(s)$ corresponds to the theta function $h^{ab}(v^a, Z^a(s)) \phi_{ab}^*\theta \left[\begin{smallmatrix} c_b^1 + D^{-1}m \\ c_b^2 \end{smallmatrix} \right] (v^b, Z^b(s))$, see Section 1.7

The set $\mathcal{B}_a := \langle \sigma_n^a, n \in \mathbb{Z}_D \rangle$ is a set of sections of $\mathcal{L}_a$ which restricts to a base of sections of $H^0(X_s, \mathcal{L}_a|_{X_s})$ for every $s \in U^{ab}$, see Section 1.5. The set $\mathcal{B}'_b := \langle \Phi_{ab}^*\sigma_m^b, m \in \mathbb{Z}_D \rangle$ is also a set of sections of $\mathcal{L}_a|_{X^{ab}}$ which restricts to a base of sections of $H^0(X_s, \mathcal{L}_a|_{X_s})$ for every $s \in U^{ab}$. There exists then a matrix $\tilde{C}^{ab}(s)$ of complex functions $U^{ab} \longrightarrow \mathbb{C}^\times$ such that $\mathcal{B}'_b = \tilde{C}^{ab}(s) \mathcal{B}_a$. Expressing this relation in terms of theta functions, it becomes

$$\begin{aligned} h^{ab}(v^a, Z^a(s)) &< \phi_{ab}^*\theta \left[\begin{smallmatrix} c_b^1 + D^{-1}m \\ c_b^2 \end{smallmatrix} \right] (v^b, Z^b(s)), m \in \mathbb{Z}_D > \\ &= \tilde{C}^{ab}(s) < \theta \left[\begin{smallmatrix} c_a^1 + D^{-1}n \\ c_a^2 \end{smallmatrix} \right] (v^a, Z^a(s)), n \in \mathbb{Z}_D >. \end{aligned}$$

By using Proposition 1.1 and relation (2) in the proof of that proposition, we conclude that $\tilde{C}^{ab}(s) = e(\pi i {}^t M[c]^1 M[c]^2 - \pi i {}^t c^1 c^2)(\det(\gamma^{ab}Z^a(s) + \delta^{ab}))^{\frac{1}{2}} C_{M^{ab}}^{-1}$ and so, $\det \tilde{C}^{ab}(s) = \zeta_8(\det(\gamma^{ab}Z^a(s) + \delta^{ab}))^{\frac{d}{2}}$, except when $3|d_g$ and $\gcd(3, d_{g-1}) = 1$, in which case we have that $\det \tilde{C}^{ab}(s) = \zeta_{24}(\det(\gamma^{ab}Z^a(s) + \delta^{ab}))^{\frac{d}{2}}$.

By the construction of the line bundle $\mathcal{L} := \mathcal{L}_{\text{can}}$ we have canonical isomorphisms $i_a : \mathcal{L}|_{X^{ab}} \xrightarrow{\cong} \mathcal{L}_a|_{X^{ab}}$ which make the following diagram commutative

$$\begin{array}{ccc} \mathcal{L}_a|_{X^{ab}} & \xrightarrow{\Phi_{ab}} & \mathcal{L}_b|_{X^{ab}} \\ & \swarrow i_a \quad \searrow i_b & \\ & \mathcal{L}|_{X^{ab}} & \end{array} \tag{6}$$

The set $\mathcal{B}_a^{\mathcal{L}} := \langle i_a^*\sigma_n^a, n \in \mathbb{Z}_D \rangle$ is a set of sections of $\mathcal{L}|_{X^a}$ which restricts to a base of sections of $H^0(X_s, \mathcal{L}|_{X_s})$ for every $s \in U^a$. Similarly, the set $\mathcal{B}_b^{\mathcal{L}} := \langle i_b^*\sigma_m^b, m \in \mathbb{Z}_D \rangle$ is a set of sections of $\mathcal{L}|_{X^b}$ which restricts to a base of sections of $H^0(X_s, \mathcal{L}|_{X_s})$ for every $s \in U^b$. According now to the above Diagram (6), we have on $X^{ab} = f^{-1}(U^{ab})$ that $\mathcal{B}_b^{\mathcal{L}} = \tilde{C}^{ab}(s) \mathcal{B}_a^{\mathcal{L}}$, where $\tilde{C}^{ab}(s)$ is the above defined matrix. To conclude, we have proven

Lemma 2.1 *Let $f : X \longrightarrow B$ be a fibration of abelian varieties of relative dimension g and $\mathcal{L}$ the above constructed symmetric (resp. totally symmetric and $g \geq 3$) line bundle on X . Fix the trivialization $\{U^a\}$ of B defined in Section 2.1. Then the transition functions of the line bundle $\det f_* \mathcal{L}$ are given by $g_{\mathcal{L}}^{ab}(s) = \zeta_8(\det(\gamma^{ab} Z^a(s) + \delta^{ab}))^{\frac{d}{2}}$ (resp. $g_{\mathcal{L}}^{ab}(s) = (\det(\gamma^{ab} Z^a(s) + \delta^{ab}))^{\frac{d}{2}}$), except when $3|d_g$ and $\gcd(3, d_{g-1}) = 1$, in which case we have that they are given by $g_{\mathcal{L}}^{ab}(s) = \zeta_{24}(\det(\gamma^{ab} Z^a(s) + \delta^{ab}))^{\frac{d}{2}}$ (resp. $g_{\mathcal{L}}^{ab}(s) = \zeta_3(\det(\gamma^{ab} Z^a(s) + \delta^{ab}))^{\frac{d}{2}}$).*

To proceed a little further we state the following lemma

Lemma 2.2 *Let $f : X \longrightarrow B$ be a fibration of abelian varieties and $s : B \longrightarrow X$ its zero section. Let $\Omega_{X/B}$ denote the relative cotangent bundle. Then $\Omega_{X/B} \cong f^* E$, where $E \cong s^* \Omega_{X/B}$ is the vector bundle on B defined by the transition matrices $g_E^{ab} := (\gamma^{ab} Z^a(s) + \delta^{ab})^{-1}$. In particular for the relative dualizing sheaf of f we have that $\omega_{X/B} \cong f^* \mu$, where $\mu \cong s^* \omega_{X/B}$ is the line bundle on B defined by the transition functions $g_\mu^{ab}(s) = \det(\gamma^{ab} Z^a(s) + \delta^{ab})^{-1}$.*

Proof. Let $\{U^a\}_{a \in A}$ be an open covering of B and $\langle U^a, \lambda_1^a(s), \dots, \lambda_{2g}^a(s) \rangle$ a choice of a symplectic base on the fibers of $X^a := f^{-1}(U^a)$, as in Section 2.1. Let $Z^a(s)$ be the period matrix of X_s^a defined by $\langle \lambda_1^a(s), \dots, \lambda_g^a(s) \rangle = \langle \lambda_{g+1}^a(s), \dots, \lambda_{2g}^a(s) \rangle Z^a(s)$. Λ_D acts on $\mathbb{C}^g \times U^a$ by $n(v, s) := (v + j_{Z^a(s)}(n), s)$, where $j_{Z^a(s)}(n) := Z^a(s)n^1 + n^2$ (where $n = (n^1, n^2)$), see Section 1.6. There is a canonical isomorphism $\phi_a : X^a \longrightarrow (\mathbb{C}^g \times U^a)/\Lambda_D$ (fibered over U^a) defined on the level of universal coverings by sending $\tilde{\phi}_a(\lambda_{g+i}^a) = (e_i, s) \in \mathbb{C}^g \times U^a$, $i = 1, \dots, g$, where $\langle e_1, \dots, e_g \rangle$ is the standard base of $\mathbb{C}^g$. Let $\langle z_1, \dots, z_g \rangle$ denote the standard coordinates of $\mathbb{C}^g$. Then dz_i is the dual to e_i . Let $z_i^a := \tilde{\phi}_a^*(z_i \times id)$. Then $\langle dz_1^a, \dots, dz_g^a \rangle$ defines at each point of X^a a base of sections of the fiber of $\Omega_{X/B}|_{X^a}$ and $dz_1^a \wedge \dots \wedge dz_g^a$ defines a (nowhere zero) section of $\omega_{X/B}|_{X^a}$. This is because $dz_1 \wedge \dots \wedge dz_g$ defines a (nowhere zero) section of the relative dualizing sheaf of the fibration $(\mathbb{C}^g \times U^a)/\Lambda_D \longrightarrow U^a$.

We have that $\langle \lambda_1^b, \dots, \lambda_{2g}^b \rangle = \langle \lambda_1^a, \dots, \lambda_{2g}^a \rangle {}^t M^{ab}$. Therefore

$$\begin{aligned} \langle \lambda_{g+1}^b, \dots, \lambda_{2g}^b \rangle &= \langle \lambda_1^a, \dots, \lambda_{2g}^a \rangle \begin{pmatrix} {}^t \gamma^{ab} \\ {}^t \delta^{ab} \end{pmatrix} = \\ &= \langle \lambda_{g+1}^a, \dots, \lambda_{2g}^a \rangle (Z^a(s), I) \begin{pmatrix} {}^t \gamma^{ab} \\ {}^t \delta^{ab} \end{pmatrix} = \langle \lambda_{g+1}^a, \dots, \lambda_{2g}^a \rangle {}^t (\gamma^{ab} Z^a(s) + \delta^{ab}). \end{aligned}$$

By taking dual bases we have that

$$\langle dz_1^b, \dots, dz_g^b \rangle = \langle dz_1^a, \dots, dz_g^a \rangle (\gamma^{ab} Z^a(s) + \delta^{ab})^{-1}$$

and so,

$$dz_1^b \wedge \dots \wedge dz_g^b = \det(\gamma^{ab} Z^a(s) + \delta^{ab})^{-1} dz_1^a \wedge \dots \wedge dz_g^a.$$

This proves the lemma. □

The proof of Theorems A and B is a consequence of Lemma 2.1 and Lemma 2.2.

2.3 The Jacobian fibration - Proof of Theorem C.

We apply now the above considerations to the Jacobian fibration $f : \mathcal{J} \longrightarrow \mathcal{M}_g$, where $\mathcal{J}$ denotes the universal Jacobian parametrizing line bundles of degree zero on the fibers of the universal curve $\psi : \mathcal{C} \longrightarrow \mathcal{M}_g$ over the moduli space $\mathcal{M}_g$ of smooth, irreducible curves of genus $g \geq 3$ without automorphisms. The Picard group of $\mathcal{M}_g$ is generated by a line bundle λ which is defined to be the determinant of the Hodge bundle of the map ψ i.e. $\lambda := \det \psi_* \omega_{\mathcal{C}/\mathcal{M}_g}$, see [1].

On the Jacobian fibration $\mathcal{J}$, there is a totally symmetric line bundle $\mathcal{L}$ which restricts to a line bundle of class 2θ on the fibers and it is trivial along the zero section. It is defined as the pull back of the Poincare bundle under the natural map. By Theorem B, we get that

Corollary 2.1 *Let $\mathcal{L}$ denote the above canonical line bundle on $\mathcal{J}$. Then*

$$\det f_*(\mathcal{L}^{\otimes n}) \cong -\frac{(2n)^g}{2} \det \psi_* \omega_{\mathcal{C}/\mathcal{M}_g}.$$

Remark 2.1 One can also prove the above Corollary 2.1 by using the Theorem 5.1 in [3]. For the case of the universal Jacobian, we can get rid from the torsion factor which appears in that theorem, by using the fact that the Picard group of $\mathcal{M}_g$ is generated by $\det \psi_* \omega_{\mathcal{C}/\mathcal{M}_g}$, see [1].

Let $f_{g-1} : \mathcal{J}^{g-1} \longrightarrow \mathcal{M}_g$ be the Jacobian fibration of degree $g-1$. We use the following result from [7], [8]. Let $\alpha : \tilde{\mathcal{M}}_g \longrightarrow \mathcal{M}_g$ denote the covering of even theta characteristics in $\mathcal{J}^{g-1}$. It is a covering of degree $2^{g-1}(2^g + 1)$. The theta divisor Θ in $\mathcal{J}^{g-1}$ intersects $\tilde{\mathcal{M}}_g$ transversely and the (set theoretic) intersection projects birationally via α to a divisor in $\mathcal{M}_g$ which has class $2^{g-3}(2^g + 1)c_1(\lambda)$. On the other hand the generic point of the intersection corresponds to a line bundle with two sections. By the description of the singularities of the

theta divisor, we have that on such a point the theta divisor has a singularity of multiplicity 2. We therefore have that the push-forward via α of the (scheme theoretic) intersection of Θ with $\tilde{\mathcal{M}}_g$ is a divisor of class $2^{g-2}(2^g + 1)c_1(\lambda)$.

We use the following commutative diagram.

$$\begin{array}{ccccccc}
 \mathcal{J} & \xleftarrow{\gamma} & \tilde{\mathcal{J}} & \xrightarrow{\phi} & \tilde{\mathcal{J}}^{g-1} & \xrightarrow{\delta} & \mathcal{J}^{g-1} \\
 \downarrow f & & \searrow \tilde{f} & & \swarrow \tilde{f}_{g-1} & & \downarrow f_{g-1} \\
 \mathcal{M}_g & \xleftarrow{\alpha} & \tilde{\mathcal{M}}_g & \xrightarrow{\alpha} & \mathcal{M}_g & &
 \end{array} \tag{7}$$

In the above diagram we denote by $\tilde{\mathcal{J}}$ and $\tilde{\mathcal{J}}^{g-1}$ the pull back of $\mathcal{J}$ and $\mathcal{J}^{g-1}$ on $\tilde{\mathcal{M}}_g$. By ϕ we denote the map which sends the line bundle $L \in \tilde{\mathcal{J}}$ sitting over the point $[C, \eta] \in \tilde{\mathcal{M}}_g$ to the line bundle $L^{\otimes 2} \otimes \eta \in \tilde{\mathcal{J}}^{g-1}$. The map ϕ is etale of degree 2^{2g} . Let $s : \tilde{\mathcal{M}}_g \rightarrow \tilde{\mathcal{J}}$ denote the zero section and let $\sigma : \tilde{\mathcal{M}}_g \rightarrow \tilde{\mathcal{J}}^{g-1}$ denote the section which sends $[C, \eta] \mapsto \eta$. We have that $\phi s = \sigma$.

Let $\tilde{\Theta}$ be the line bundle corresponding to the theta divisor on $\tilde{\mathcal{J}}^{g-1}$. Then $\tilde{\Theta} = \delta^* \Theta$ and so, $\alpha_* c_1(\tilde{f}_{g-1}^* \tilde{\Theta}^{\otimes n}) = 2^{g-1}(2^g + 1)c_1(f_{g-1}^* \Theta^{\otimes n})$. Let $\tilde{\lambda}$ be the determinant of the Hodge bundle of the fibration $\tilde{f}$. Then $\tilde{\lambda} = \alpha^* \lambda$ and so $\alpha_* c_1(\tilde{\lambda}) = 2^{g-1}(2^g + 1)c_1(\lambda)$. From the things we stated above, we have that if $\tilde{\mu} := \sigma^* \tilde{\Theta}$ then $\alpha_* c_1(\tilde{\mu}) = 2^{g-2}(2^g + 1)c_1(\lambda)$. Let $\tilde{\mathcal{L}}$ be the canonical line bundle on $\tilde{\mathcal{J}}$ of Corollary 2.1. Then $\tilde{\mathcal{L}} = \gamma^* \mathcal{L}$.

One can see that $c_1(\tilde{f}_* \phi^* \tilde{\Theta}^{\otimes n}) = 2^{2g} c_1(\tilde{f}_{g-1}^* \tilde{\Theta}^{\otimes n})$. A proof can be given by applying the GRR theorem. One can also see that the restriction of $\phi^* \tilde{\Theta}$ on a fiber of the map $\tilde{f}$ is equal to the restriction of $\tilde{\mathcal{L}}^{\otimes 2}$ on the same fiber. This can be seen by using Proposition 3.5 of Ch.2 in [4] and the Riemann's constant theorem. Therefore, by the see-saw principle, the line bundles $\tilde{\mathcal{L}}^{\otimes 2}$ and $\phi^* \tilde{\Theta}$ are isomorphic up to tensor by a line bundle which is pull back of a line bundle from $\tilde{\mathcal{M}}_g$. Since $s^* \tilde{\mathcal{L}}^{\otimes 2} \cong \mathcal{O}$ and $s^* \phi^* \tilde{\Theta} \cong \tilde{\mu}$, we have that

$$\tilde{\mathcal{L}}^{\otimes 2n} \otimes \tilde{f}^* \tilde{\mu}^{\otimes n} \cong \phi^* \tilde{\Theta}^{\otimes n}.$$

By applying $\tilde{f}_*$ and taking c_1 we have that

$$c_1(\tilde{f}_* \tilde{\mathcal{L}}^{\otimes 2n}) + (4n)^g n c_1(\tilde{\mu}) = 2^{2g} c_1(\tilde{f}_{g-1}^* \tilde{\Theta}^{\otimes n}).$$

Apply now α_* to get

$$-2^{g-1}(2n)^g 2^{g-1}(2^g + 1)c_1(\lambda) + (4n)^g n 2^{g-2}(2^g + 1)c_1(\lambda) = 2^{2g} 2^{g-1}(2^g + 1)c_1(f_{g-1}^* \Theta^{\otimes n}).$$

Therefore

$$c_1(f_{g-1} * \Theta^{\otimes n}) = \frac{1}{2} n^g (n-1) c_1(\lambda).$$

By using the fact that the $\text{Pic} \mathcal{M}_g$ is generated by λ , see [1], this concludes the proof of the Theorem B.

2.4 Alternative proof of Theorem C.

We give now the alternative proof of Theorem C promised in the introduction. This is an application of the GRR theorem, see also Appendice 2 in [5] for a similar calculation. We keep the notation of the above section 2.3. In the above Diagram 7 let ϕ be the map which sends the line bundle $L \in \tilde{\mathcal{J}}$ sitting over the point $[C, \eta] \in \tilde{\mathcal{M}}_g$ to the line bundle $L \otimes \eta \in \tilde{\mathcal{J}}^{g-1}$. By Lemma 2.2 we have that $\Omega_{\tilde{\mathcal{J}}/\tilde{\mathcal{M}}_g} \cong \tilde{f}^* \tilde{s}^* \Omega_{\tilde{\mathcal{J}}/\tilde{\mathcal{M}}_g}$ and since ϕ is an isomorphism, we get that $\Omega_{\tilde{\mathcal{J}}^{g-1}/\tilde{\mathcal{M}}_g} \cong \tilde{f}_{g-1}^* \tilde{s}^* \Omega_{\tilde{\mathcal{J}}^{g-1}/\tilde{\mathcal{M}}_g}$.

We proceed now to the proof of Theorem C: We apply GRR to the fibration $\tilde{f}_{g-1} : \tilde{\mathcal{J}}^{g-1} \longrightarrow \tilde{\mathcal{M}}_g$. It is

$$\text{ch}(\tilde{f}_{g-1}!(\tilde{\Theta}^{\otimes n})) = \tilde{f}_{g-1}*(\text{ch}(\tilde{\Theta}^{\otimes n}) \cdot \text{td}(\Omega_{\tilde{\mathcal{J}}^{g-1}/\tilde{\mathcal{M}}_g}^\vee)).$$

We get that

$$c_1(\tilde{f}_{g-1} * \tilde{\Theta}^{\otimes n}) = \frac{n^{g+1}}{(g+1)!} f_* c_1^{g+1}(\tilde{\Theta}) - \frac{n^g}{2} c_1(\tilde{\lambda}).$$

The vanishing of the terms on the right hand side containing the “factor” c_1^k with $k \leq g-1$ in the expansion of $\text{ch}(\tilde{\Theta}^{\otimes n})$, is a consequence of the projection formula and the fact that $\Omega_{\tilde{\mathcal{J}}^{g-1}/\tilde{\mathcal{M}}_g} \cong \tilde{f}_{g-1}^* E$, where E is a vector bundle, see Lemma 2.2. The form of the term containing the “factor” c_1^{g-1} is due to the Poincare formula. The appearance of $\tilde{\lambda}$ is a consequence of Corollary 2.1.

Lets say now that $c_1(f_{g-1} * \Theta^{\otimes n}) = a(n) c_1(\lambda)$ and that $f_* c_1^{g+1}(\Theta) = b c_1(\lambda)$, where $a(n), b \in \mathbb{Z}$, see [1]. Then $c_1(\tilde{f}_{g-1} * \tilde{\Theta}^{\otimes n}) = a(n) c_1(\tilde{\lambda})$ and $\tilde{f}_* c_1^{g+1}(\Theta) = b c_1(\tilde{\lambda})$. We get $a(n) = \frac{n^{g+1}}{(g+1)!} b - \frac{n^g}{2}$. For $n = 1$ the above gives that $b = (g+1)!(a(1) + \frac{1}{2})$. Now $a(1) = 0$, because the line bundle $\tilde{f}_{g-1} * \tilde{\Theta}$ has by definition a nowhere zero section, so it is the trivial bundle. We then get $b = \frac{(g+1)!}{2}$ and so, $c_1(f_{g-1} * \Theta^{\otimes n}) = \frac{1}{2} n^g (n-1) c_1(\lambda)$.

References

- [1] Arbarello E. and Cornalba M.: *The Picard group of the moduli spaces of curves*. Topology 26, 152-171 (1987).

- [2] Brasch H-J.: *Lifting level D -structures of abelian varieties*. Arch. Math. 60, 553-562 (1993).
- [3] Faltings G. and Chai Ch-L.: *Degeneration of Abelian Varieties*. Springer-Verlag, 1990.
- [4] Lange H. and Birkenhake Ch.: *Complex Abelian Varieties*. Springer-Verlag, 1992.
- [5] Moret-Bailly, L.: *Pinceaux de varietes abeliennes*. Asterisque 129 (1985).
- [6] Mumford D.: *Tata Lectures on Theta I*. Progress in Mathematics, vol. 28, Birkhäuser 1983.
- [7] Teixidor M.: *Half-canonical series on algebraic curves*. Transactions of A.M.S. 302, 99-115 (1987).
- [8] Teixidor M.: *The divisor of curves with a vanishing theta-null*. Comp. Math. 66, 15-22 (1988).